

Exclusive Central Hadron Production in $p\bar{p}$ Collisions at the Tevatron for $\sqrt{s} = 1960\text{GeV}, 900\text{GeV}$

Partial Wave Analysis - Full Status

M. Żurek, A. Świąch
Jagiellonian University, Kraków

D. Lontkovskyi, I. Makarenko
University of Kyiv

M. Albrow, J.S. Wilson, J. Lewis
FNAL



Contents

Physics Motivation

Data sample

Gap cuts

Exclusivity cuts

2 Exclusive tracks on primary vertex

- Additional cuts
- Acceptance calculation
- Systematic uncertainties
- Partial Wave Analysis



Physics Motivation

Double Pomeron Exchange



Pomeron:

- Carrier of 4-momentum between protons
- Strongly interacting color singlet combination of quarks or/and gluons
- Quantum numbers of vacuum
- LO: $P = gg$

Single Diffraction



Double Pomeron Exchange



Analysis

GXG reaction

$$\bar{p} + p \rightarrow \bar{p} + \text{GAP} + X + \text{GAP} + p$$

X (in this study):

- hadron pair mostly $\pi^+ \pi^-$
- central $y \approx 0$
- between rapidity gaps $\Delta y \approx 4$
- $Q = S = 0$, $C = +1$, $J = 0$ or 2 , $I=0$

Expected to be dominated by DPE in the t-channel!

Low Mass Central Hadronic State Analysis

Candidates selection

Trigger requirement:

- 2 central ($|\eta| < 1.3$) towers with $E_t > 0.5$ GeV
- PCAL ($2.11 < |\eta| < 3.64$) in veto
- CLC ($3.75 < |\eta| < 4.75$) in veto
- BSC1 ($5.4 < |\eta| < 5.9$) in veto

Gap cuts:

To determine noise levels in subdetectors we divide zero-bias sample from same periods into two sub-samples:

No Interaction:

- No tracks and
- No CLC hits and
- No muon stubs

Interaction:

At least one


- Track or
- CLC hit or
- Muon stub

Low Mass Central Hadronic State Analysis


Candidates selection

Examples of exclusive requirements – empty forward detectors

CDF Run II Preliminary, $\sqrt{s}=1960\text{GeV}$



CDF Run II Preliminary, $\sqrt{s}=1960\text{GeV}$




Low Mass Central Hadronic State Analysis

Candidates selection

Exclusivity cuts

To determine exclusive 2-4 tracks we apply similar technique in central region, just excluding cones of $R=0.3$ around each track extrapolation.

$$R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$$



Effective exclusive luminosity

- Determination of efficiency of having no-pileup using zero-bias sample.

We measure ratio of empty events (all detectors on noise level) to all events.

- Exponential drop with bunch luminosity.
- Slope corresponds to total detected inelastic cross section.

1960 GeV

900 GeV


53.88(36) mb

62.76(38) mb


1.18/pb

0.059/pb

CDF Run II Preliminary



CDF Run II Preliminary



Low Mass Central Hadronic State Analysis

Candidates selection


Additional cuts:

- quality of tracks
 ΔZ_0 , N of COT hits, χ^2/N_{dof} , p_T
- cosmic ray rejection:
no muons, 3D opening angle, d_0
- Physical cuts:
 η , rapidity of central state, total charge


Examples:

d_0 , $y(X)$

CDF Run II Preliminary, $\sqrt{s}=1960$ GeV



CDF Run II Preliminary, $\sqrt{s}=1960$ GeV



Cut	1960GeV	900GeV
Triggered	92130 x 10 ³	21737 x 10 ³
Forward cleanup	59276 x 10 ³	18749 x 10 ³
2 tracks	4700 x 10 ³	271 x 10 ³
Quality + cosmic rejection + exclusivity	415413	8400
Opposite sign	350009	7595
Luminosity	7.12/pb	0.074/pb
Exclusive luminosity	1.18/pb	0.059/pb

Acceptance calculation

Model independent analysis 3 components:

Kinematics cuts:

- $P_t(\pi) > 0.4 \text{ GeV}/c$
- $|\eta(\pi)| < 1.3$
- $|y(\pi)| < 1.0$
- Trigger efficiency
- Single track acceptance
- 2 tracks acceptance

Trigger efficiency

1. Sample of min-bias data, good quality isolated (no other tracks in cone with $R=0.4$) tracks.
2. Checking how often they fired 0, 1, 2 or more trigger towers (≥ 4 bits) in 3×3 box around track extrapolation.
3. Trigger efficiency composed from those 3 probability distributions (which are functions of P_t and η)

Before: trigger efficiency only in the function of P_t

Trigger efficiency

Probability of triggering 2 or more towers in the central detector by two independent tracks „a” and „b”:

$$\varepsilon = P_2(a) + P_1(a) * [P_1(b) + P_2(b)] + P_0(a) * P_2(b)$$

P_0 – probability of triggering no towers




P_1 – probability of triggering one tower

P_2 – probability of triggering two or more towers

Trigger efficiency


	$P_2(b)$	$P_1(b)$	$P_0(b)$
$P_2(a)$	X	X	X
$P_1(a)$	X	X	
$P_0(a)$	X		

Trigger efficiency




Single track acceptance

1. Single pion generation, flat in ϕ
2. Acceptance in the function of $P_t(\text{track})$ and η
 - Probability that track will be reconstructed at all
 - Probability that track will pass all single track quality cuts




Single track acceptance


Single track acceptance in η




Single track acceptance in P_t



Single track acceptance in P_t and η



Single track acceptance




2 tracks cuts acceptance

Cuts:

- 3D opening angle
- y of central state
- Separation
- dZ0

Based on J=0 phase space model

All previous cuts applied before







Systematical uncertainties

1. All cut parameters varied by
 - ~ 1 standard deviation (gaussian-like) or
 - ~ 0.5 of FWHM width (Lorentz like)what looks reasonably (others)
2. Trigger efficiency – statistical uncertainties of probability distributions
3. Same value of cut for E-W forward detectors.
4. Assumed independence of such systematics.
5. Applied simultaneously in data/MC


Pt distribution for different mass ranges

rebless



Mass distribution – tail fit

rebless



Question 1

Do we want to present all our spectra from $0.28\text{GeV}/c^2$ in mass or from $0.8\text{GeV}/c^2$?

Partial Waves Analysis - Idea

Unpolarized coss-section

$$\begin{aligned} \frac{d\sigma}{d\Omega} = & \frac{1}{(2s_a + 1)(2s_b + 1)p^2} \sum_{(\lambda), J, J'} \left(J + \frac{1}{2}\right) \left(J' + \frac{1}{2}\right) (-1)^{\lambda - \mu} \\ & \cdot \langle \lambda_a \lambda_b | T_J(E) | \lambda_c \lambda_d \rangle * \langle \lambda_a \lambda_b | T_{J'}(E) | \lambda_c \lambda_d \rangle \\ & \cdot \sum_l C(JJ'l; \lambda, -\lambda) C(JJ'l; \mu, -\mu) P_l(\cos \theta) \end{aligned}$$

M.Jacob, G.C.Wick, On the general theory of collisions for particles with spin, Ann. Phys. **7**, (1959) 404-428.

$$a + b \rightarrow c + d$$

- ▶ s_a, s_b - spins
- ▶ J, J' - total angular momenta
- ▶ $\lambda_a, \lambda_b, \lambda_c, \lambda_d$ - helicities; $\mu = \lambda_c - \lambda_d, \lambda = \lambda_a - \lambda_b$
- ▶ p - momentum of initial state particle, E - c.m. energy
- ▶ $T = i(1 - S)$, S - scattering matrix
- ▶ $C(JJ'l; \lambda, -\lambda)$ - C-G coefficients

Double Pomeron Exchange

$$\text{Goal: } \langle \lambda_a \lambda_b | T_J(E) | \lambda_c \lambda_d \rangle = ?$$

DPE properties:

- ▶ $\pi^+ \pi^-$ production only via s -channel diagrams
- ▶ $0^{++}, 2^{++}, 4^{++}, \dots$ intermediate states only
 - each such state has a definite J
 - 0^{++} states contribute only to T_0
- ▶ $s_\pi = 0, s_{\mathbb{P}} = 0, \lambda_\pi, \lambda_{\mathbb{P}} = 0$
- ▶ Therefore: $\langle \lambda_a \lambda_b | T_J(E) | \lambda_c \lambda_d \rangle$ is a single complex number $\mathbf{R}_J(\mathbf{E}) e^{i\phi_J(\mathbf{E})}$

Tool: Measurement of coefficients of Legendre polynomials a_l

0^{++} and 2^{++} central state assumption

- ▶ $J, J' = 0, 2 \rightarrow I = 0, 2, 4$.
- ▶ Only non-zero C-G coefficients: $C(000; 00)$, $C(022; 00)$, $C(220; 00)$, $C(222; 00)$, $C(224; 00)$

1. $I = 4 \rightarrow$ only $J = J' = 2$:

$$a_4 = \left(\frac{9}{7}\right)^2 p^{-2} R_2^2$$

2. $I = 0 \rightarrow J = J' = 2$ or $J = J' = 0$:

$$a_0 = \frac{1}{4} p^{-2} (R_0^2 + R_2^2)$$

3. $I = 2 \rightarrow J = J' = 2$ or $J = 0, J' = 2$ or $J = 2, J' = 0$:

$$a_2 = p^{-2} \left(\frac{5}{2} R_0 R_2 \cos(\phi_2 - \phi_0) + \left(\frac{5}{7}\right)^2 R_2^2 \right),$$

where: $\delta = \phi_2 - \phi_0$ - relative phase

Finally:

1. $R_2^2 = p^2 \left(\frac{7}{9}\right)^2 a_4$

2. $R_0^2 = p^2 \left(4a_0 - \left(\frac{7}{9}\right)^2 a_4 \right)$

3. $\delta = \frac{1}{2} \frac{a_2 - \left(\frac{5}{9}\right)^2 a_4}{\sqrt{\left(\frac{7}{9}\right)^2 a_4 \left(4a_0 - \left(\frac{7}{9}\right)^2 a_4 \right)}}$

Legendre moments - correction for acceptance

Corrected Legendre moments **b** (vector):

$$\mathbf{b} = \mathbf{K}^{-1} \mathbf{a}.$$

a - not corrected Legendre moments, **K** following matrix:

$$K_{ll'} = \frac{\sum_i w_i P_l(\cos \theta_i) P_{l'}(\cos \theta_i)}{\sum_i w_i},$$

where: $w_i = w_i^{MC} \cdot w_i^{Acc}$.

<http://www-cdf.fnal.gov/jsw/internal/GXG/PWA-corrections.md.html>

Legendre moments - correction for acceptance

1. Statistical uncertainties:

$$\text{cov}(b_l, b_{l'}) = K_{ll'}^{-1} \text{cov}(a_l, a_{l'}) \left(K_{ll'}^{-1} \right)^T$$

We need the covariance of the mean value of the sample.

$$\begin{aligned} \text{cov}(a_l, a_{l'}) &= \frac{\sum_{ij} w_i w_j \text{cov}(P_l(\cos \theta_i), P_{l'}(\cos \theta_j))}{\sum_{ij} w_i w_j} \\ &= \frac{\sum_i w_i^2}{\sum_{ij} w_i w_j} \text{cov}(P_l(\cos \theta), P_{l'}(\cos \theta)) \end{aligned}$$

Let us denote: $V_1 = \sum_i w_i$, $V_2 = \sum_i w_i^2$, then:

$$\begin{aligned} \text{cov}(a_l, a_{l'}) &= \frac{V_2}{V_1^2} \frac{V_1}{V_1^2 - V_2} \sum_i w_i (P_l(\cos \theta_i) - a_l)(P_{l'}(\cos \theta_i) - a_{l'}) \\ &= \frac{V_2}{V_1^2 - V_2} \left(\frac{\sum_i w_i P_l P_{l'}}{V_1} - a_l a_{l'} \right) = \frac{V_2}{V_1^2 - V_2} (\langle P_l P_{l'} \rangle - a_l a_{l'}) \end{aligned}$$

Legendre moments - correction for acceptance

2. Uncertainties linked with K^{-1} matrix: related to statistics of our MC sample

M. Lefebvre, R.K. Keeler, R. Sobie, J. White, Propagation of Errors for Matrix Inversion, [arXiv:hep-ex/9909031]

Let us denote: $\epsilon_{lm} = \langle P_l P_m \rangle$:

$$\text{cov}(\epsilon_{ab}^{-1}, \epsilon_{cd}^{-1}) = \sum_{ijkl} \epsilon_{ai}^{-1} \epsilon_{jb}^{-1} \epsilon_{ck}^{-1} \epsilon_{ld}^{-1} \text{cov}(\epsilon_{ij}, \epsilon_{kl}),$$





thus:

$$\delta b_i^2 = \sum_{jk} a_j \text{cov}(\epsilon_{ab}^{-1}, \epsilon_{cd}^{-1}) a_k$$





$\text{cov}(\epsilon_{ab}^{-1}, \epsilon_{cd}^{-1})$ - calculated in analogous way as in 1.

3. Systematical uncertainties: We varied all parameters (in Data and MC) and checked the result in Legendre moments plots.




MC – no weighting







MC – no weighting







MC – no weighting






Data – no MC weighting







Data – no MC weighting







Data – no MC weighting






MC – weighting







MC – weighting







MC – weighting






Data – MC weighting



Data – MC weighting



Data – MC weighting



Conclusions

- **We have measured $\pi^+\pi^-$ pairs between large rapidity gaps at the Tevatron, which should be dominated by double pomeron exchange. The background from K^+K^- is small.**
- **We do not see a $\rho(770)$, confirming that photoproduction and ρ -exchange, are negligible.**
- **This is the only measurement from the Tevatron, and has much higher statistics than preliminary data from the LHC experiments.**
- **The mass spectra show several structures:**
 - Broad continuum below $1 \text{ GeV}/c^2$,
 - Sharp drop at $1 \text{ GeV}/c^2$
 - Resonant enhancement around $1.0 - 1.5 \text{ GeV}/c^2$.
- **The s-dependence is mass dependent.**
- **We plan to do a partial wave analysis to distinguish different spin states.**

Thank you

Backup slides

Data sample

- Datasets used:
 - gdifap – 1960 GeV
 - gdifar – 900/300 GeV
- Same trigger requirement:
DIFF_TWO_CJET0.5_PLUGVETO_0.75
 - 2 central ($|\eta| < 1.32$) towers with $E_t > 0.5$ GeV
 - Plug ($2.11 < |\eta| < 3.64$) in veto ($E_t < 0.75$ GeV)
 - BSC1 and CLC in veto
- $L = 7.12/\text{pb}$ – 1960 GeV and $L = 0.074/\text{pb}$ – 900 GeV
- Gaps at least from $|\eta| = 1.3$ to $|\eta| = 5.9$

Effective exclusive luminosity

Efficiency of having no-pileup using zero-bias sample. We measure ratio of empty events (all detectors on noise level) to all events.

Should drop exponentially with bunch luminosity and be equal 1 at $L=0$


Slope corresponds to part of inelastic cross section with particles in $|\eta| < 5.9$. (More low mass diffraction is included at 900 GeV than at 1960 GeV.):

- 53.88(36) mb – 1960 GeV
- 62.76(38) mb – 900 GeV


• Effective luminosities:

- 1.18/pb – 1960 GeV
- 0.059/pb – 900 GeV

CDF Run II Preliminary



CDF Run II Preliminary



Systematics - summary

Cut	1960 GeV	900 GeV
Exclusivity cut in CD	15%	15%
Forward Plug	6%	6%
Pt	2%	2%
Z0	2%	4%
BSC	2%	4%
Luminosity	6%	6%
χ^2	3%	3%
Total:	20%	20%